



## 64. SAVETOVANJE 64<sup>th</sup> CONFERENCE

# PROIZVODNJA I PRERADA ULJARICA

sa međunarodnim učešćem

# PRODUCTION AND PROCESSING OF OILSEEDS

with international participation

## ZBORNIK RADOVA PROCEEDINGS

Herceg Novi, Crna Gora  
25 - 30. jun 2023. godine



## IZDAVAČI PUBLISHERS

UNIVERZITET U NOVOM SADU, TEHNOLOŠKI FAKULTET NOVI SAD  
UNIVERSITY OF NOVI SAD, FACULTY OF TECHNOLOGY NOVI SAD  
INSTITUT ZA RATARSTVO I POVRTARSTVO NOVI SAD,  
INSTITUT OD NACIONALNOG ZNAČAJA ZA REPUBLIKU SRBIJU  
INSTITUTE OF FIELD AND VEGETABLE CROPS NOVI SAD,  
NATIONAL INSTITUTE OF THE REPUBLIC OF SERBIA  
„INDUSTRIJSKO BILJE“ DOO NOVI SAD  
„INDUSTRIAL PLANTS“ DOO NOVI SAD

## UREĐIVAČKI ODBOR EDITORIAL BOARD

Prof. dr Biljana Pajin, Prof. dr Ranko Romanić, Dr Vladimir Miklič, Dr Vojin Đukić  
Mr Zvonimir Sakač, Dr Olga Čurović, Zoran Nikolovski, dipl. inž., Vladimir Šarac,  
dipl. inž., Gordan Parenta, dipl. inž., Nada Grbić, dipl. inž., Milan Ševo, dipl. inž.,  
Dragan Trzin, dipl. inž.

## UREDNIK EDITOR Savet tehnologa

## TEHNIČKI UREDNICI TECHNICAL EDITORS

Prof. dr Ranko Romanić  
Doc. dr Ivana Lončarević

## ADRESA IZDAVAČA PUBLISHER'S ADDRESS

„INDUSTRIJSKO BILJE“ DOO, NOVI SAD  
21000 Novi Sad, Dimitrija Tucovića 2A, Srbija  
Tel/fax. +381 21 66 16 633, +381 21 66 24 311, +381 21 66 12 135  
e-mail: office@indbilje.co.rs

ISBN 978-86-6253-170-4

## ŠTAMPA PRINT



Štamparija Feljton, Novi Sad  
Stražilovska 17  
Tel: 021/ 66-22-867





Jelena Rabrenović, Milica Fotirić Akšić, Aleksandra Rašović, Dragana Dabić Zagoračić, Milica Sredojević, Ivanka Čirić, Nataša Obradović, Mina Volić, Maja Natić	VALORIZACIJA SEMENA MALINE U CILJU DOBIJANJA HLADNO PRESOVANOG ULJA I BIOAKTIVNIH EKSTRAKATA IZ POGAČE VALORIZATION OF RASPBERRY SEEDS IN ORDER TO OBTAIN COLD-PRESSED OIL AND BIOACTIVE EXTRACTS FROM OIL CAKE .....	129
Jovana Pantić, Senka Popović, Danijela Šuput, Nevena Hromiš, Ljiljana Popović, Ranko Romanić	ANTIOKSIDATIVNI POTENCIJAL BIOPOLIMERNIH FILMOVA NA BAZI POGAČE SEMENA ŠLJIVE ANTIOXIDATIVE POTENTIAL OF BIOPOLYMER FILMS BASED ON PLUM SEED CAKE .....	141
Vesna Vujasinović, Bojan Đerčan, Milan Vukić, Dragan Vujadinović, Dajana Bjelajac, Goran Radivojević, Danijela Rajić, Kristina Šarenac	CHIA SEME: DA LI JE ZAISTA SUPERHRANA SA ASPEKTA SASTAVA MASNIH KISELINA? CHIA SEEDS: IS IT REALLY A SUPERFOOD FROM THE ASPECT OF FATTY ACID COMPOSITION? .....	149
György Karlovits	STRATEGIJA RAZVOJA NOVOG JESTIVOGL ULJA ZA GENERACIJU SENIORA STRATEGY FOR THE DEVELOPMENT OF A NEW EDIBLE OIL FOR THE SENIOR GENERATION .....	157
Petar Ilić, Vojislav Banjac, Olivera Đuragić, Slađana Rakita, Bojana Kokić, Viktor Stojkov, Ana Marjanović Jeromela	MOGUĆNOST UPOTREBE HLADNO CEĐENOG ULJA LANIKA U ISHRANI KUĆNIH LJUBIMACA THE POSSIBILITY OF APPLICATION OF COLD PRESSED CAMELINA SEED OIL IN PET FOOD .....	159
Gordan Parenta, Ranko Romanić, Tanja Lužaić, Petar Klać, Marija Gvozdenović, Branislav Milković, Milivoj Števanov, Stevan Švenderman, Nenad Vlahović	UTICAJ FILTRACIJE I KLARIFIKACIJE NA KVALITET SIROVOG PRESOVANOG SUNOKRETOVOG ULJA INFLUENCE OF FILTRATION AND CLARIFICATION ON THE QUALITY OF CRUDE PRESSED SUNFLOWER OIL .....	167



# MOGUĆNOST UPOTREBE HLADNO CEĐENOGLJULJA LANIKA U ISHRANI KUĆNIH LJUBIMACA

Petar Ilić<sup>1</sup>, Vojislav Banjac<sup>1</sup>, Olivera Đuragić<sup>1</sup>, Slađana Rakita<sup>1</sup>,  
Bojana Kokić<sup>1</sup>, Viktor Stojkov<sup>1</sup>, Ana Marjanović Jeromela<sup>2</sup>

<sup>1</sup>Univerzitet u Novom Sadu,

Naučni institut za prehrambene tehnologije u Novom Sadu, Novi Sad, Srbija

<sup>2</sup>Institut za ratarstvo i povrтарstvo, Institut od nacionalnog značaja za  
Republiku Srbiju, Novi Sad, Srbija

## IZVOD

Cilj ovog istraživanja je ispitivanje nutritivne vrednosti i mogućnost primene hladno ceđenog ulja lanika dobijenog iz semena dva srpska genotipa lanika (NS Zlatka i NS Slatka) u hrani za kućne ljubimce. U tu svrhu određivan je masnokiselinski sastav, odnos  $\omega$ -6 i  $\omega$ -3 masnih kiselina, kao i aterogeni i trombogeni indeksi i hipoholesterolski/hiperholesterolski odnos. Takođe, određivan je i sadržaj liposolubilnih vitamina, bioaktivnih komponenti i antioksidativni kapacitet. Prikazani su i rezultati analize boje.

**Ključne reči:** uljarice, hladno ceđeno ulje, hrana za kućne ljubimce,  $\omega$ -3 masne kiseline, tokoferoli

## THE POSSIBILITY OF APPLICATION OF COLD PRESSED CAMELINA SEED OIL IN PET FOOD

### ABSTRACT

The aim of this research is to investigate the nutritive value and the possibility of application of cold pressed camelina seed oil obtained from two Serbian camelina seed genotypes (NS Zlatka and NS Slatka) as a potential ingredient for pet food. For that purpose fatty acid profil,  $\omega$ -6 and  $\omega$ -3 ratio, atherogenic, thrombogenicity indexes and ratio of hypocholesterolemic to hypercholesterolemic were analysed. The study also included the investigation of the content of fat-soluble vitamins and the content of bioactive compounds as well as antioxidant capacity. The color of cold pressed oil was also determined.

**Key words:** oilseed, cold pressed oil, pet food,  $\omega$ -3 fatty acids, tocopherol



## UVOD

Lanik (*Camelina sativa* L.) je uljana biljna vrsta iz porodice kupusnjača (Brassicaceae), koja se uzgajala još 4000 godina p.n.e. Kultivacija lanika se vršila sporadično u Evropi sve do sredine 20. veka kada biva zamenjen produktivnijim vrstama kao što su suncokret i uljana repica. (Zubr, 2010; Berti i sar., 2016; Juodka i sar., 2022). Lanik je tek poslednjih desetak godina počeo da dobija sve veću pažnju naučne zajednice zbog brojnih mogućnosti iskorišćenja, od proizvodnje biogoriva, materijala i agrohemikalija do prehrambene industrije i industrije hrane za životinje (Minkowski i sar., 2011). Takođe, prednosti uzgoja lanika su te što poseduje kratak period vegetacije i niske zahteve za đubrivotom i vodom. Prilagodljiv je na različite uslove okoline, toleriše hladnu klimu i suše, i u poređenju sa ostalim uljaricama može da se uzgaja na marginalnom i zasoljenom tlu. Zbog sadržaja pojedinih fitohemikalija, lanik pokazuje i otpornost na štetočine i pojedine izazivače bolesti, što omogućava smanjenu upotrebu pesticida i herbicida (Mondor i Hernández-Álvarez, 2022).

Omega-6 ( $\omega$ -6) i omega ( $\omega$ -3) masne kiseline su esencijalne u ishrani pasa, jer psi nemaju sposobnost njihove sinteze. Poželjan odnos  $\omega$ -6 i  $\omega$ -3 masnih kiselina za pse iznosi 5:1 i 10:1, što ulje lanika, koje je veoma bogato  $\omega$ -3 masnim kiselinama, čini idealnim za podešavanje ovog odnosa u hrani (Vaughh i sar., 2021). Dokazano je da  $\omega$ -3 masne kiseline kod pasa imaju kardioprotективни efekat (Billman i sar., 1997), kao i antiinflamatornu i imunomodulatornu ulogu (Wander i sar., 1997). Takođe, unapređuju izgled i zdravlje kože i krvna (Kirby i sar., 2009).

Seme lanika u proseku sadrži 24-35 % proteina i 36 % ulja koje sadrži 40-60% polinezasićenih masnih kiselina, od čega  $\alpha$ -linoleinska kiselina čini 35-40 % (Raczyk i sar., 2016; Juodka i sar., 2022). Za određivanje nutritivne vrednosti hladno ceđenih ulja, pored masnokiselinskog sastava bitan je i sadržaj tokoferola, sterola, karotenoidnih pigmenata i hlorofila (Ratusz i sar., 2018). Seme lanika je bogato antioksidantima, posebno tokoferolima, fitosterolima, fenolnim kiselinama i flavonoidima (Zanetti i sar., 2021; Mondor i Hernández-Álvarez, 2022). Tokoferoli, kao prirodni antioksidanti, imaju ključnu ulogu u prevenciji oksidacije polinezasićenih masnih kiselina u ulju (Ratusz i sar., 2018). Visok sadržaj  $\omega$ -3 masnih kiselina i tokoferola, čini ulje lanika pogodnom sirovinom za proizvodnju hrane za pse (Burron i sar., 2021). Prirodno prisutni pigmenti u ulju su važni zbog toga što mogu da daju nepoželjnu tamnu boju, a mogu i da doprinesu oksidaciji (Ratusz i sar., 2018). Hladno ceđeno ulje lanika je zlatno žute boje čija nijansa zavisi od prisustva i sadržaja materija kao što su karotenoidi i hlorofili (Sizova i sar., 2003).

Genotipi lanika korišćeni u ovom istraživanju su razvijeni u Srbiji kako bi se proširio portfolio dostupnih vrsta u balkanskoj agrikulturi. Ovi genotipi poseduju dobre proizvodne karakteristike i razvijeni su da budu prilagođeni klimatskim uslovima koji vladaju na Balkanu (Marjanović Jeromela i sar., 2021; Čanak et al., 2022).



## MATERIJAL I METODE RADA

U eksperimentalnom radu korišćeno je hladno ceđeno ulje dva srpska genotipa lanika (*Camelina sativa* L.), NS Zlatka i NS Slatka, koji su razvijeni na Institutu za ratarstvo i povrtarstvo u Novom Sadu i koji su registrovani kod Ministarstva poljoprivrede, šumarstva i vodoprivrede Republike Srbije 2018. godine.

Sadržaj masnih kiselina analiziran je na gasnom hromatografu Agilent 7890A (Agilent Technologies, Santa Clara, CA, USA) gde je korišćena kapilarna kolona sa silicijum-dioksidom SP SP-2560 (100 m × 0.25 mm, d=0.20 µm; Supelco, Bellefonte, USA) Priprema uzorka izvršena je prema metodi SRPS EN ISO 12966-2:2017.

Aterogeni (AI) i trombogeni (TI) indeksi izračunati su primenom formule koju je predložio Ulbricht i Southgate (1991), dok je hipoholesterolski / hiperholesterolski odnos izračunat primenom formule predložene od strane autora Santos i sar. (2002).

Tokoferoli su analizirani po metodi SRPS EN 12822:2014 gde je korišćen HPLC-FLD (Agilent 1260 Infinity, Agilent Technologies, Santa Clara, CA, USA) sa kolonom sa normalnom fazom (Phenomenex Luna Silica, 5 µm, 250 mm x 4.6 mm). Rezultati su izraženi u mg tokoferola po 100 g suvog uzorka (mg/ 100g). Vitamin A je određivan po metodi SRPS EN 12823-1:2014, dok je vitamin D3 određivan po metodi SRPS EN 12821:2008.

Sadržaj β-karotena je određen prema metodi koju su primenili Savić Gajić i sar. (2019). Sadržaj ukupnih polifenola određen je primenom UV- VIS spektrofotometrije i Folin-Ciocalteau reagensa. Antioksidativni potencijal određen je primenom DPPH testa (Savić Gajić i sar., 2019) i procenjena je na osnovu IC<sub>50</sub> vrednosti koja je dobijena interpolacijom.

Boja uzoraka instrumentalno je određena upotrebom Minolta Chroma Meter CR 400 (Konica Minolta Sensing Inc., Japan). Rezultati su prezentovani prema CIELab sistemu boja, gde su koordinate definisane na sledeći način: L\* je koordinata svetloće boje (0 označava crnu, a 100 belu boju), a\* je crveno-zelena koordinata (a\*+ označava crvenu, a a\*- označava zelenu boju) i b\* je žuto-plava koordinata (b\*+ označava žutu, a b\*- označava plavu boju).

## REZULTATI I DISKUSIJA

Masnokiselinski sastav hladno ceđenog ulja lanika prikazan je u Tabeli 1. Analiza masnih kiselina pokazala je da hladno ceđeno ulje lanika poseduje nutritivno vredan sastav sa najvišim procentom polinezasićenih masnih kiselina (PUFA) (~58 %), sadržaja mononezasićenih masnih kiselina oko 32 % i zasićenih masnih kiselina oko 9 %. Od zasićenih masnih kiselina (SFA) najveći udio zauzima palmitinska kiselina (C16:0), dok su miristinska, stearinska i arahidinska kiselina pokazale manji sadržaj. Od mononezasićenih masnih kiselina (MUFA) najdominantnije su gondoinska (~15%) i oleinska kiselina (~13%), dok je erukinska kiselina bila prisutna u manjoj meri (~3%). Dobijeni rezultati su uporedivi sa rezultatima dobijenim u istraživanjima koje su sproveli Budin i sar. (1995), Eidhin (2003) i Zubr i sar. (2002).



Tabela 1. Sadržaj masnih kiselina u hladno ceđenom ulju lanika  
Table 1. The fatty acid content of cold pressed camelina seed oil

Masne kiseline (%) Fatty acids (%)	NS Zlatka	NS Slatka
C14:0	0,04	0,05
C16:0	5,39	5,35
C16:1	0,10	0,10
C18:0	2,50	2,48
C18:1n9c	13,40	13,87
C18:2n6c	15,89	15,83
C20:0	1,30	1,18
C20:1n9	15,44	15,17
C18:3n3	38,76	39,25
C20:2n6	2,11	2,03
C22:1n9	3,18	2,91
C20:3n3	1,89	1,77
Zasićene masne kiseline (SFA)	9,23	9,07
Mononezasićene masne kiseline (MUFA)	32,12	32,05
Polinezasićene masne kiseline (PUFA)	58,65	58,88
ω-3	40,65	41,02
ω-6	18,00	17,86
ω-6/ω-3	0,44	0,44
Aterogeni indeks (AI)	0,06	0,06
Trombogeni indeks (TI)	0,05	0,05
Hipoholesterolski/hiperholesterolski odnos FA (HH)	12,54	12,75

Takođe, analiza je pokazala da je hladno ceđeno ulje lanika izuzetno bogato n-3 i n-6 polinezasićenim masnim kiselinama, od kojih je najprisutnija α-linoleinska kiselina (~39%), dok je udeo linolne kiseline iznosio oko 16%. Ove masne kiseline su esencijalne jer psi nemaju sposobnost za njihovu sintezu. Sadržaj α-linoleinske kiseline u hladno ceđenom ulju lanika je veći nego kod soje, suncokreta, konoplje i uljane repice, ali manji nego kod lana (Čolović i sar., 2015; Zajac i sar., 2020; Joudka i sar., 2022). Odnos n-6 i n-3 masnih kiselina je takođe bitan faktor u evaluaciji nutritivne vrednosti ulja. U ovom istraživanju taj odnos je iznosio 0.44, što ovo ulje čini pogodnim za primenu u proizvodnji hrane za kućne ljubimce (Burron i sar., 2021).

Za procenu nutritivne vrednosti ulja, određivani su i aterogeni (AI) i trombogeni (TI) indeksi, kao i hipoholesterolski/ hiperholesterolski odnos (HH). Ove vrednosti bolje opisuju nutritivnu vrednost ulja nego masnokiselinski profil (Ratusz i sar., 2018). Aterogeni i trombogeni indeksi bili su znatno niži od 1, dok su HH indeksi bili relativno visoki 12,54 - 12,75, što je u skladu sa preporučenim vrednostima sa nutritivnog aspekta kvaliteta ulja.



Tabela 2. Sadržaj liposolubilnih vitamina u hladno ceđenom ulju lanika  
Table 2. The content of fat-soluble vitamins of the cold pressed camelina seed oil

Vitamin (mg/100 g)	NS Zlatka	NS Slatka
Vitamin (mg/100 g)		
α-tokoferol	2,11	2,68
β-tokoferol	<0,1	<0,1
γ-tokoferol	38,51	54,74
A	<0,1	<0,1
D3	<0,02	<0,02

Sadržaj tokoferola u srpskim genotipima lanika prikazan je u tabeli 2. Najprisutniji tokoferol u hladno ceđenom ulju genotipa NS Zlatka i NS Slatka jeste γ-tokoferol (38,51 mg/100g i 54,74 mg/100g, respektivno), dok je α-tokoferol bio prisutan u manjoj meri (2,11 mg/100g i 2,68 mg/100g, respektivno), što je uporedivo sa rezultatima istraživanja Ratusz i sar. (2018). U istraživanju koje su sproveli Grajzer i sar. (2020), dobijene vrednosti sadržaja tokoferola su bile nešto veće i iznosile su 817,7 mg/kg za γ-tokoferol, dok je sadržaj α-tokoferola bio ispod granica detekcije. U ovom istraživanju sadržaji β-tokoferola, vitamina A i D3 bili su ispod granica detekcije kod oba genotipa.

Tabela 3. Sadržaj bioaktivnih jedinjenja i antioksidativni potencijal  
hladno ceđenog ulja lanika  
Table 3. The content of bioactive compounds and antioxidative potential  
of cold pressed camelina seed oil

Bioaktivna jedinjenja i antioksidativni potencijal	NS Zlatka	NS Slatka
β-karoten (mg/kg)	89,43	71,12
Total polyphenols (mg GKE/100 g)	7,3	30,9
IC <sub>50</sub> [mg/mL]	9,4	9,5

Sadržaj β-karotena prikazan u tabeli 3 kod genotipa NS Zlatka i NS Slatka iznosio je 89,43 mg/kg i 71,12 mg/kg, respektivno, što je uporedivo sa rezultatima istraživanja Raczyk i sar. (2016), gde je dobijen sadržaj β-karotena iznosio 78 to 112 mg/kg. Sadržaj ukupnih polifenola kod NS Zlatke iznosio je 7,3 mg GKE/100g, dok je kod NS Slatke bio nešto viši i iznosio je 30,9 mg GKE/100 g. U istraživanju koje su sproveli Ergönül i Özbek (2018), sadržaj ukupnih polifenola u hladno ceđenim uljima različitih vrsta lanika varirao je između 25,9 i 112,7 mg GAE/kg, što je niža vrednost u poređenju sa dobijenim rezultatima u ovom istraživanju. Ulje oba genotipa je pokazalo dobar antioksidativni potencijal, gde je IC<sub>50</sub> vrednost bila približno ista i iznosila je 9,4 mg/ml za NS Zlatku i 9,5 mg/ml za NS Slatku.




Tabela 4. Boja hladno ceđenog ulja lanika  
Table 4. The color of cold pressed camelina seed oil

Parametar	NS Zlatka	NS Slatka
L*	59,98	61,82
a*	-6,06	-6,63
b*	55,97	50,27

Rezultati određivanja CIE L\*a\*b\* koordinata hladno ceđenog ulja semena lanika dati su u tabeli 4. Hladno ceđeno ulje semena lanika genotipa NS Zlatka je prema parametru L\* bilo nešto tamnije od genotipa NS Slatka, dok je sudeći po parametru b\* bilo žuće. Ulje NS Slatke je imalo nešto nižu vrednost parametra a\*, što ga čini „zelenijim“ u odnosu na ulje NS Zlatke.

## ZAKLJUČAK

Dobijeni rezultati ovog istraživanja ukazuju na to da je hladno ceđeno ulje semena lanika genotipa NS Zlatka i NS Slatka bogat izvor esencijalnih masnih kiselina i tokoferola, što ga čini pogodnom alternativom za upotrebu u proizvodnji hrane za pse. Takođe, utvrđene niske vrednosti aterogenog i trombogenog indeksa, kao i relativno visoka vrednost hipoholesterolsko/hiperholesterolskog odnosa ukazuju na to da hrana obogaćena ovim uljem može da smanji učestalost oboljevanja od koronarnih bolesti kod pasa.

## Zahvalnica

Ovaj rad je finansijski podržalo Ministarstvo Nauke, Tehnološkog Razvoja i Inovacija Republike Srbije (451-03-47/2023-01/200222).

## LITERATURA

1. Berti, M., Gesch, R., Eynck, C., Anderson, J., Cermak, S. (2016). Camelina uses, genetics, genomics, production, and management. Industrial Crops and Products, 94, 690-710.
2. Billman, G. E., Kang, J. X., Leaf, A. (1997). Prevention of ischemia-induced cardiac sudden death by n-3 polyunsaturated fatty acids in dogs. Lipids, 32(11), 1161-1168.
3. Budin, J. T., Breene, W. M., & Putnam, D. H. (1995). Some compositional properties of camelina (*Camelina sativa* L. Crantz) seeds and oils. Journal of the American Oil Chemists' Society, 72, 309-315.
4. Burron, S., Richards, T., Patterson, K., Grant, C., Akhtar, N., Trevizan, L., Pearson, W., Shoveller, A. K. (2021). Safety of Dietary Camelina Oil Supplementation in Healthy, Adult Dogs. Animals, 11(9), 2603.
5. Čanak, P., Zanetti, F., Jovičić, D., Vujošević, B., Miladinov, Z., Stanisavljević, D., Miroslavljević, M., Alberghini, B., Facciolla, E., Jeromela, A. M. (2022). Camelina



- guation under osmotic stress –Trend lines, time-courses and critical point. Industrial Crops and Products, 181, 114761.
6. Čolović, D. S., Berenji, J. J., Levart, A. R., Salobir, J. K., Lević, J. D., Čolović, I., & Spović, S. J. (2015). Fatty acid composition and natural antioxidant capacity of ten Serbian linseed cultivars. Food and Feed Research, 42(2), 139-145.
7. Eidhin, D. N., Burke, J., & O'beirne, D. (2003). Oxidative stability of  $\omega$ 3-rich camelina oil and camelina oil-based spread compared with plant and fish oils and sunflower spread. Journal of Food Science, 68(1), 345-353.J.
8. Grajzer, M., Szmalcel, K., Kuźmiński, Ł., Witkowski, M., Kulma, A., & Prescha, A. (2020). Characteristics and antioxidant potential of cold-pressed oils - Possible strategies to improve oil stability. Foods, 9(11), 1630.
9. Günç Ergönül, P., Aksoylu Özbek, Z. (2018). Identification of bioactive compounds and total phenol contents of cold pressed oils from safflower and camelina seeds. Journal of Food Measurement and Characterization, 12, 2313-2323.
10. Juodka, R., Nainienė, R., Juškienė, V., Juška, R., Leikus, R., Kadžienė, G., Stankevičienė, D. (2022). Camelina (*Camelina sativa* (L.) Crantz) as feedstuffs in meat type poultry diet: A source of protein and n-3 fatty acids. Animals, 12(3), 295.
11. Kirby, N. A., Hester, S. L., Rees, C. A., Kennis, R. A., Zoran, D. L., Bauer, J. E. (2009). Skin surface lipids and skin and hair coat condition in dogs fed increased total fat diets containing polyunsaturated fatty acids. Journal of animal physiology and animal nutrition, 93(4), 505-511.
12. Marjanović Jeromela, A., Cvejić, S., Mladenov, V., Kuzmanović, B., Adamović, B., Stojanović, D., Vollmann, J. (2021). Technological quality traits phenotyping of Camelina across multienvironment trials. Acta Agriculturae Scandinavica, Section B—Soil & Plant Science, 71(8), 667-673.
13. Minkowski, K., Grzeskiewicz, S., Jerzewska, M., Rolno-Spozywczego, I. B. P. (2011). Assessment of nutritive value of plant oils with high content of linolenic acids based on the composition of fatty acids, tocopherols, and sterols. *Zywnosc Nauka Technologia Jakosc* (Poland).
14. Mondor, M., Hernández-Álvarez, A. J. (2022). Camelina sativa composition, attributes, and applications: A review. European Journal of Lipid Science and Technology, 124(3), 2100035.
15. Raczyk, M., Popis, E., Kruszewski, B., Ratusz, K., & Rudzińska, M. (2016). Physicochemical quality and oxidative stability of linseed (*Linum usitatissimum*) and camelina (*Camelina sativa*) cold-pressed oils from retail outlets. European Journal of Lipid Science and Technology, 118(5), 834-839.
16. Ratusz, K., Symoniuk, E., Wroniak, M., Rudzińska, M. (2018). Bioactive compounds, nutritional quality and oxidative stability of cold-pressed camelina (*Camelina sativa* L.) oils. Applied Sciences, 8(12), 2606.
17. Santos-Silva, J., Bessa, R. J. B., Santos-Silva, F. J. L. P. S. (2002). Effect of genotype, feeding system and slaughter weight on the quality of light lambs: II. Fatty acid composition of meat. Livestock Production Science, 77(2-3), 187-194.
18. Savić, I., Savić Gajić, I., Gajić, D. (2020). Physico-chemical properties and oxidative stability of fixed oil from plum seeds (*Prunus domestica* Linn.). Biomolecules, 10(2), 294.



- Chikunova, N. V., Pikulev, I. V., Chikunova, T. M. (2003). Fatty acid composition of *Camelina sativa* L. Crantz oil and the selection of an optimal antioxidant. Khimično-tekhnicheskoj promstvo i nafta, 19(2), 27-31.
- PS EN 12821:2008. Srpski standard. Prehrambeni proizvodi - Određivanje vitamina D tečnom hromatografijom visoke performanse - Merenje holekalciferola D3 i ergokalciferola D2, Institut za standardizaciju Srbije, Beograd.
21. SRPS EN 12822:2014. Srpski standard. Prehrambeni proizvodi- Određivanje vitamina E tečnom hromatografijom visoke performanse - Merenje  $\alpha$ ,  $\beta$ ,  $\gamma$  i  $\delta$  tokoferola, Institut za standardizaciju Srbije, Beograd.
22. SRPS EN 12823-1:2014. Srpski standard. Prehrambeni proizvodi - Određivanje vitamina A tečnom hromatografijom visoke performanse - Deo 1: Merenje svih E-retinola i 13-Z-retinola, Institut za standardizaciju Srbije, Beograd.
23. SRPS EN ISO 12966-2:2017. Srpski standard. Ulja i masti biljnog i životinjskog porekla - Gasna hromatografija metilestara masnih kiselina - Deo 2: Priprema metilestara masnih kiselina, Institut za standardizaciju Srbije, Beograd.
24. Ulbricht, T. L. V., Southgate, D. A. T. (1991). Coronary heart disease: seven dietary factors. *The lancet*, 338(8773), 985-992.
25. Vaughn, D. M., Reinhart, G. A., Swaim, S. F., Lauten, S. D., Garner, C. A., Boudreaux, M. K., Spano, J. S., Hoffman, C.E., Conner, B. (1994). Evaluation of effects of dietary n-6 to n-3 fatty acid ratios on leukotriene B synthesis in dog skin and neutrophils. *Veterinary Dermatology*, 5(4), 163-173.
26. Wander, R. C., Hall, J. A., Gradin, J. L., Du, S. H., Jewell, D. E. (1997). The ratio of dietary (n-6) to (n-3) fatty acids influences immune system function, eicosanoid metabolism, lipid peroxidation and vitamin E status in aged dogs. *The Journal of nutrition*, 127(6), 1198-1205.
27. Zajac, M., Kiczorowska, B., Samolińska, W., Klebaniuk, R. (2020). Inclusion of camelina, flax, and sunflower seeds in the diets for broiler chickens: Apparent digestibility of nutrients, growth performance, health status, and carcass and meat quality traits. *Animals*, 10(2), 321.
28. Zanetti, F., Alberghini, B., Marjanović Jeromela, A., Grahovac, N., Rajković, D., Kiprovska, B., Monti, A. (2021). Camelina, an ancient oilseed crop actively contributing to the rural renaissance in Europe. A review. *Agronomy for Sustainable Development*, 41(1), 1-18.
29. Zubr, J. (2010). Carbohydrates, vitamins and minerals of *Camelina sativa* seed. *Nutrition & Food Science*, 40(5), 523-531.
30. Zubr, J., Matthäus, B. (2002). Effects of growth conditions on fatty acids and tocopherols in *Camelina sativa* oil. *Industrial crops and products*, 15(2), 155-162.



СИР - Каталогизација у публикацији  
Библиотеке Матице српске, Нови Сад

633.85(082)  
665.3(082)

**САВЕТОВАЊЕ “Производња и прерада уљарица” (64 ; Херцег Нови ; 2023)**

Zbornik radova / 64. savetovanje “Proizvodnja i prerada uljarica” sa međunarodnim učešćem = Proceedings / 64th Conference “Production and Processing of Oilseeds” with international participation, Herceg Novi, 25 - 30. jun 2023. - Novi Sad : Tehnološki fakultet : Institut za ratarstvo i povrtarstvo : Industrijsko bilje, 2023 (Novi Sad : Feljton). - 263 str. : ilustr. ; 25 cm

Tiraž 150. - Rezime na engl. jeziku uz svaki rad. - Bibliografija uz svaki rad. - Registar.

ISBN 978-86-6253-170-4

a) Уљарице - Производња - Зборници б) Уљарице - Прерада - Зборници

COBISS.SR-ID 117401865